

Percolation on sparse random graphs with given degree sequence

N. Fountoulakis*
 School of Mathematics
 University of Birmingham
 Edgbaston, B15 2TT
 United Kingdom
 nikolaos@maths.bham.ac.uk

February 2, 2008

Abstract

We study the two most common types of percolation process on a sparse random graph with a given degree sequence. Namely, we examine first a bond percolation process where the edges of the graph are retained with probability p and afterwards we focus on site percolation where the vertices are retained with probability p . We establish critical values for p above which a giant component emerges in both cases. Moreover, we show that in fact these coincide. As a special case, our results apply to power law random graphs. We obtain rigorous proofs for formulas derived by several physicists for such graphs.

1 Introduction

Traditionally percolation theory has been the study of the properties of a random subgraph of an infinite graph, that is obtained by deleting each edge of the graph with probability $1 - p$ for some $p \in (0, 1)$ independently of every other edge. The question that has been mainly investigated is whether the subgraph that is spanned by these edges has an infinite component or not. The classical type of graphs that was studied in percolation theory is the lattice $\mathbb{Z}^d$ in various dimensions $d \geq 2$ (see [24]). Until now various other types of lattices have been studied. In each of the above cases the main problem is the calculation of a critical p_c so that if $p < p_c$ then the random subgraph obtained as above has no infinite components, whereas if $p > p_c$ there is an infinite component with probability 1.

In the present work, we study percolation on finite graphs whose number of vertices is large. This problem is old, in the sense that for example a $\mathcal{G}_{n,p}$ random graph is a random subgraph of the complete graph on n vertices, where each edge appears with probability p independently of every other edge. In this context, a question about the appearance of an infinite component is senseless. A somehow analogous question is whether there exists a component of the random subgraph containing a certain proportion of the vertices or as we customarily say a *giant component*. More specifically, if the original graph has n vertices the question now is whether there exists an $\varepsilon > 0$ for which there is a component of the random subgraph that has at least εn vertices with probability $1 - o(1)$ (as $n \rightarrow \infty$). Hence, we also ask (quite informally) for the existence of a critical p_c for which whenever $p < (1 - \delta)p_c$ then for every $\varepsilon > 0$ there is

*The author is supported by the EPSRC, grant no. EP/D50564X/1

no component having at least εn vertices with probability $1 - o(1)$ and whenever $p > p_c(1 + \delta)$ then there is a component with at least εn vertices for some $\varepsilon > 0$ with probability $1 - o(1)$. A classical example of this is the $\mathcal{G}_{n,p}$ random subgraph of K_n , the complete graph on n vertices, where as it was proved by Erdős and Rényi in [20] the critical probability is equal to $1/n$ (see also [8] or [25] for an extensive discussion).

More generally, Bollobás, Kohayakawa and Łuczak in [11] raised the following question: given a sequence of graphs $\{G_n\}$ whose order tends to infinity as n grows, is there such a phase transition? Assume that G_n has $|G_n|$ vertices and e_n edges. For each such n we have a probability space on the set of spanning subgraphs of G_n and the probability of such a subgraph of G_n that has e edges is $p^e(1-p)^{e_n-e}$, where e_n is the number of edges of G_n . Let $G_n(p)$ be a sample from this probability space. Thus we are seeking a p_c such that: if $p < (1-\delta)p_c$, then for every $\varepsilon > 0$ as $n \rightarrow \infty$ all the components of $G_n(p)$ have at most $\varepsilon|G_n|$ vertices with probability $1 - o(1)$, and if $p > p_c(1 + \delta)$, then there exists $\varepsilon = \varepsilon(p) > 0$ for which the largest component of $G_n(p)$ has at least $\varepsilon|G_n|$ vertices with probability $1 - o(1)$. If the sequence of graphs is $\{K_n\}$, this is simply the case of a $\mathcal{G}_{n,p}$ random graph.

Other families of sequences have also been studied. For example, percolation on the hypercube with 2^n vertices has been analysed by Ajtai, Komlós and Szemerédi in [1] where the critical edge probability turns out to be also equal to $1/n$. More detailed analysis of this phase transition was carried out recently by Borgs, Chayes, van der Hofstad, Slade and Spencer in [15].

On the other hand recent research has also focused on finite graphs with bounded maximum degree. Here we consider sequences of graphs $\{G_n\}$, where each G_n is a graph on n vertices, with uniformly bounded maximum degree. Alon, Benjamini and Stacey have investigated percolation on such sequences of graphs in [4]. Among other things, they proved that the critical probability for the emergence of a component of linear size in a d -regular graph on n vertices whose girth tends to infinity with n is $1/(d-1)$ (Theorem 3.2 in [4]). Phase transitions on specific sequences of finite graphs were studied more closely by Borgs, Chayes, van der Hofstad, Slade and Spencer in [13], [14].

More recently, in [10], Bollobás, Borgs, Chayes and Riordan analysed the phase transition in sequences of dense graphs that are convergent in a certain sense.

Also, in [22] Frieze, Krivelevich and Martin, proved that $p_c = 1/d$ for sequences of d -regular graphs on n vertices which are quasi-random, when $d \rightarrow \infty$ as n grows. These are graphs whose structure resembles that of a d -regular random graph.

In the present paper, we determine a percolation threshold in the case where the sequence $\{G_n\}_{n \in \mathbb{Z}^+}$ is a sequence of sparse random graphs on n vertices. In particular, for every integer $n \geq 1$, G_n is a uniformly random graph on the set $V_n = \{1, \dots, n\}$ having a given degree sequence $\mathbf{d}(n) = (d_1, \dots, d_n)$, i.e. for $i = 1, \dots, n$ vertex i has degree d_i . More formally, a *degree sequence* on the set V_n is a vector $\mathbf{d} = (d_1, \dots, d_n)$ consisting of natural numbers, where $d_1 \leq \dots \leq d_n$, and $\sum_{i=1}^n d_i$ is even. We let $2M$ denote this sum, and $M = M(n)$ is the number of edges that $\mathbf{d}$ spans. For a given $\mathbf{d} = \mathbf{d}(n)$, if $\mathbf{d}(n) = (d_1, \dots, d_n)$ for $n \in \mathbb{Z}^+$, we set $D_i = D_i(n) = |\{j \in V_n : d_j = i\}|$, for $i \in \mathbb{N}$ and $\Delta = \Delta(n) = \max_{1 \leq i \leq n} \{d_i\} = d_n$. Finally, if G is a graph on V_n , then $D(G)$ denotes its degree sequence.

An *asymptotic degree sequence* is a sequence $(\mathbf{d}(n))_{n \in \mathbb{Z}^+}$, where for each $n \in \mathbb{Z}^+$ the vector $\mathbf{d}(n)$ is a degree sequence on V_n . An asymptotic degree sequence is *sparse*, if for every $i \in \mathbb{N}$, we have $\lim_{n \rightarrow \infty} D_i(n)/n = \lambda_i$, for some $\lambda_i \in [0, 1]$, where $\sum_{i \geq 0} \lambda_i = 1$, and moreover

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i \geq 1} i(i-2)D_i(n) = \sum_{i \geq 1} i(i-2)\lambda_i < \infty. \quad (1)$$

This implies that for every $\varepsilon > 0$ there exists $i^*(\varepsilon)$ and $N = N(\varepsilon)$ such that for every $n > N$ we have

$$\left| \frac{1}{n} \sum_{i \leq i^*} i(i-2)D_i(n) - \sum_{i \geq 1} i(i-2)\lambda_i \right| < \varepsilon. \quad (2)$$

The *generating polynomial* of a sparse asymptotic degree sequence is defined as $L(s) = \sum_{i=0}^{\infty} \lambda_i s^i$. We assume that every asymptotic degree sequence $(\mathbf{d}(n))_{n \in \mathbb{Z}^+}$ we work with is such that for every n the set of simple graphs that have $\mathbf{d}(n)$ as their degree sequence is non-empty.

We consider two types of percolation. Firstly, for some $p \in (0, 1)$, each edge of G_n is present with probability p independently of every other edge. This type of percolation is usually called *bond percolation*, in that we randomly delete the edges (i.e. the bonds) of G_n . This is distinguished from another type of percolation which is called *site percolation*. Here, we go through the vertices of G_n and we make each of them isolated with probability $1 - p$, independently of every other vertex (or as we say we delete this vertex). The random subgraph in this case is the spanning subgraph of G_n that does not contain the edges that are attached to the vertices that were deleted. The terms “bond” and “site” percolation have their origins in the percolation theory of infinite graphs (see [24] for an extensive discussion on both types as well as the references therein).

We shall now define the percolation threshold in each of the above cases. Let $G'(n)$ denote the random subgraph that is obtained in either case and let $L_1(G'(n))$ be the lexicographically first component of $G'(n)$ (this is the component that has maximum order and the smallest vertex it contains is smaller than the smallest vertex of every other component of maximum order - the comparison between the vertices is by means of the total ordering on V_n). Starting from the bond percolation we set $p_c^{\text{bond}} = \sup\{p \in [0, 1] : |L_1(G'(n))|/n \xrightarrow{p} 0 \text{ as } n \rightarrow \infty\}$ (the symbol $\xrightarrow{p}$ denotes convergence in probability, i.e. we say that $X_n \xrightarrow{p} 0$ if for every $\varepsilon > 0$ we have $\mathbb{P}[|X_n| > \varepsilon] \rightarrow 0$ as $n \rightarrow \infty$). The convergence in probability is meant with respect to the sequence of probability spaces indexed by the set $\mathbb{Z}^+$, where for each $n \in \mathbb{Z}^+$ the probability of a certain spanning subgraph is the probability that this is the subgraph which is spanned by the edges that survive the random deletion of the edges of the random graph G_n . Similarly, in the case of site percolation we define $p_c^{\text{site}} = \sup\{p \in [0, 1] : |L_1(G'(n))|/n \xrightarrow{p} 0 \text{ as } n \rightarrow \infty\}$, where $G'(n)$ is now the spanning subgraph of G_n that is the outcome of the deletion of those edges that attached to the chosen vertices, i.e. the vertices that we make isolated. Note that in both cases there are two levels of randomness.

If G_n is a random d -regular graph on V_n , for any fixed $d \geq 3$, the bond percolation threshold has been calculated by Goerdts in [23] and is equal to $1/(d-1)$. Before this, bond percolation in random regular graphs was studied by Nikolettseas, Paley, Spirakis and Yung in [30], where it was proved that the critical probability is at most $32/d$, for d large enough. Also, Nikolettseas and Spirakis in [29] study the edge expansion properties of the giant component that remains after the edge deletion process. However, these papers did not provide any analysis on the site percolation process. Our main theorem involves also the latter and is stated as follows:

Theorem 1.1 *If $(\mathbf{d}(n))_{n \in \mathbb{Z}^+}$ is a sparse asymptotic degree sequence of maximum degree $\Delta(n) \leq n^{1/9}$ and $L(s)$ is its generating polynomial which is twice differentiable at 1 and moreover $L''(1) > L'(1)$, then $p_c^{\text{site}} = p_c^{\text{bond}} = L'(1)/L''(1)$. Moreover, whenever $p > p_c^{\text{bond}}$ ($p > p_c^{\text{site}}$, respectively) there exists an $\varepsilon > 0$ such that $|L_1(G'(n))| > \varepsilon n$ with probability $1 - o(1)$.*

The formula for both critical probabilities was obtained by Dorogovtsev and Mendes in [19] using qualitative (i.e. non-rigorous) arguments.

To make the statement of the above theorem slightly clearer, let us consider the case of bond percolation (the case of site percolation is similar). Let $\mathcal{G}(n)$ be the set of graphs on V_n whose degree sequence is $\mathbf{d}(n)$. Each graph $G \in \mathcal{G}(n)$ gives rise to a probability space which consists of all its spanning subgraphs. In particular, if G has e edges and G' is a spanning subgraph of G that has $e' \leq e$ edges then its probability is $p^{e'}(1-p)^{e-e'}$; let $\mathbb{P}_p^G[\cdot]$ denote this measure. In other words, this space accommodates the outcomes of the bond percolation process applied to G and we call it the *percolation space* of G . For any $\varepsilon \in (0, 1)$ we let $g_\varepsilon(G)$ be the set of all spanning subgraphs of G whose largest component has at least εn vertices. This event has probability $\mathbb{P}_p^G[g_\varepsilon(G)]$ in the percolation space of G . Now, assume that $p < p_c^{\text{bond}}$. Theorem 1.1 implies that for any given $\rho \in (0, 1)$, the event $\{G \in \mathcal{G}(n) : \mathbb{P}_p^G(g_\varepsilon(G)) < \rho\}$ occurs with probability $1 - o(1)$ in the uniform space $\mathcal{G}(n)$. That is, asymptotically for almost every graph in $\mathcal{G}(n)$ the random deletion of the edges leaves a component of order at least εn with probability no more than ρ . If $p > p_c^{\text{bond}}$, then the second part of the theorem implies that there exists $\varepsilon > 0$ such that the event $\{G \in \mathcal{G}(n) : \mathbb{P}_p^G(g_\varepsilon(G)) > 1 - \rho\}$ occurs with probability $1 - o(1)$ in $\mathcal{G}(n)$. Hence, as $n \rightarrow \infty$ almost all the graphs in $\mathcal{G}(n)$ are such that if we apply the bond percolation process to them with retainment probability p , then there is a component having at least εn vertices with probability at least $1 - \rho$ (in the percolation space).

The fact that the critical probabilities coincide reflects a behaviour that is similar to that of percolation on an infinite regular tree. Of course in that context the critical probabilities are defined with respect to the appearance of an infinite component that contains the (vertex that has been selected as the) root. Using the fundamental theorem of Galton-Watson processes (see for example [5]), it can be shown that the bond and the site critical probabilities coincide and they are equal to $1/(d-1)$, where d is the degree of each vertex of the tree. Observe that for the case of a random d -regular graph the above theorem implies that $p_c^{\text{site}} = p_c^{\text{bond}} = 1/(d-1)$. This is not a coincidence as it is well-known that a random d -regular graph locally (e.g. at distance no more than i from a given vertex for some fixed i) looks like a d -regular tree.

More generally, the typical local structure of the class of random graphs we are investigating are also tree-like. Note that the ratio $L''(1)/L'(1)$ equals

$$\sum_{i=2}^{\infty} (i-1) \frac{i\lambda_i}{\sum_{j=1}^{\infty} j\lambda_j}. \quad (3)$$

Consider a vertex $v \in V_n$ which has positive degree and let us examine more closely the behaviour of one of its neighbours. It can be shown that the probability that this has degree i is proportional to $iD_i(n)$. In particular, it is almost equal to $\frac{iD_i(n)}{\sum_i iD_i(n)}$ and this tends to $\frac{i\lambda_i}{\sum_{j=1}^{\infty} j\lambda_j}$ as n grows. Moreover, one can show that with probability $1 - o(1)$ there are no edges between the neighbours of v . Therefore (3) is the limit of the expected number of children a neighbour of v has. This scenario is repeated for every vertex in the d -th neighbourhood of v , where d is fixed. More precisely, the vertices which are at distance no more than d induce a tree rooted at v which contains at most $\ln \ln n$ vertices, with probability $1 - o(1)$. Suppose that there are t_i vertices of degree i in this tree. Thus for a vertex that is at distance d from v , the probability that it has degree i is proportional to $i(D_i(n) - t_i) = i\lambda_i n(1 - o(1))$. More precisely, it is

$$\frac{i(D_i(n) - t_i)}{\sum_i i(D_i(n) - t_i)}.$$

Since $\Delta \leq n^{1/9}$ and $t_i \leq \ln \ln n$, it follows that $\sum_i it_i \leq \ln \ln n \sum_{i \leq n^{1/9}} i = O(n^{1/3})$. Hence, the limit of the above probability as $n \rightarrow \infty$ is again $\frac{i\lambda_i}{\sum_{j=1}^{\infty} j\lambda_j}$ and (3) gives the limiting expected

number of children of such a vertex. In other words, the graph that is induced by the vertices which are at distance no more than d from v behaves like the tree of a branching process that started at v , with the ratio $L''(1)/L'(1)$ being the expected progeny of each vertex. Observe here that the condition $L''(1) > L'(1)$ implies that in fact this is a supercritical branching process which yields an infinite tree with probability 1.

Therefore, at least locally either bond or site percolation is essentially percolation on such a random rooted tree. In both types of percolation, if $p < L'(1)/L''(1)$, then the expected number of children of a vertex that survive is $pL''(1)/L'(1) < 1$. Thus the random tree that is developing around v after the random failures of the edges or the vertices will be distributed as the tree of a subcritical branching process. In particular, the tree that surrounds most of the vertices will be cut off from the rest of the graph at a relatively small depth. On the other hand, if $p > L'(1)/L''(1)$ a large proportion from each of these local trees is preserved and moreover they are big enough to guarantee that there are enough edges going out of them. So eventually there is a fair chance that some of them are joined together and form a component of linear order. However, this is only a qualitative approach to Theorem 1.1. The actual proof and the structure of the paper are described in Section 2.

1.1 Theorem 1.1 and power-law graphs

The power-law degree sequences are those for which for any $k \geq 1$ one has $\lambda_k = ck^{-\gamma}$, for some constants $c, \gamma > 0$. We should point out that the crucial parameter here is γ . Such degree sequences have attracted much attention in the last few years mainly because of the fact that they arise in “natural” networks such as the Internet, the WWW or biological networks (see [12], [2] or [19] for a survey of results or the recent book by Chung and Lu [17] for a more detailed discussion). For example, in [21] Faloutsos, Faloutsos and Faloutsos gave evidence that the Internet as it looked like in 1995, viewed as a graph whose vertices are the routers, and the edges are the physical links between them, has a power-law degree sequence with $\gamma \approx 2.48$. Bond and site percolation on such networks naturally correspond to random failures of the links or of the nodes, respectively. Thus, a site percolation process on the Internet may be seen as random failures of routers.

For a power-law degree sequence with $\gamma > 3$ one has $L(s) = c \sum_{k \geq 2} \frac{s^k}{k^\gamma}$ where $|s| \leq 1$. Thus, if $\zeta(\lambda) = \sum_{k \geq 1} \frac{1}{k^\lambda}$ is the Riemann’s zeta function, then $L'(1) = c \sum_{k \geq 2} \frac{k}{k^\gamma} = c \sum_{k \geq 2} \frac{1}{k^{\gamma-1}} = c\zeta(\gamma-1)$ and $L''(1) = c \sum_{k \geq 2} \frac{k(k-1)}{k^\gamma} = c \sum_{k \geq 2} \frac{1}{k^{\gamma-2}} - c \sum_{k \geq 2} \frac{1}{k^{\gamma-1}} = c(\zeta(\gamma-2) - \zeta(\gamma-1))$ (of course here the derivatives are left derivatives). Let $\gamma_0 = \sup\{\gamma : \gamma > 3, \frac{\zeta(\gamma-2)}{\zeta(\gamma-1)} > 2\}$. Theorem 1.1 implies that the critical probabilities for a power-law degree sequence with $3 < \gamma < \gamma_0$ but maximum degree at most $n^{1/9}$ are

$$p_c^{site} = p_c^{bond} = \frac{\zeta(\gamma-1)}{\zeta(\gamma-2) - \zeta(\gamma-1)}. \quad (4)$$

This agrees with the analysis made in [16] by Callaway, Newman, Strogatz and Watts for the case of site percolation on a random graph whose degree sequence follows a “truncated” power-law, that is $\lambda_k = Ck^{-\gamma}e^{-k/\kappa}$, for $C, \kappa > 0$. As $\kappa \rightarrow \infty$, then this approaches a power-law distribution with parameter γ . It can be shown that in this case the critical probability they obtain converges to the expression in (4) (see for example Equation (141) p.45 in [2]). Similar analysis made by Cohen, Erez, ben-Avraham and Havlin in [18] suggests that if $\gamma \leq 3$ there is no phase transition at all. Also, Boguñá, Pastor-Satorras and Vespignani in [7] argue that this happens whenever $2 < \gamma \leq 3$. This was also suggested by simulations in [3]. In particular, Albert, Jeong and

Barabási give experimental evidence of the result of a site percolation process on a random graph (obtained from a different model) whose degree sequence is power-law with $\gamma = 3$. They observe that the graph remains largely intact by the random deletion of vertices and no threshold behaviour is observed. That is, even if a large proportion of vertices are deleted, there is always a component of linear order. In the same paper, the authors give also experimental evidence in samples of the Internet and the World-Wide Web, concluding that no phase transition occurs even for small values of p . However, Dorogovtsev and Mendes in [19] applying the formula of Theorem 1.1 (which they also obtain in their paper), but without our degree restrictions, give the scaling of the critical probabilities as functions of n as $n \rightarrow \infty$, for $2 < \gamma \leq 3$. In that context the critical probabilities are defined empirically, according to whether or not the proportion of vertices in the largest component after the percolation process is almost zero.

The case $\gamma \leq 3$ corresponds to $L''(1)$ being divergent which suggests that p_c^{site} and p_c^{bond} vanish. However, Theorem 1.1 works under the assumption that $L''(1)$ converges. It would be an interesting and natural next step to prove or disprove the existence of a positive critical probability in the case where $L''(1)$ is divergent.

2 Definitions and sketch of the proof

In this paper we are interested in sparse asymptotic degree sequences $\mathcal{D}$ satisfying $Q(\mathcal{D}) := \sum_{i=1}^{\infty} i(i-2)\lambda_i > 0$. This is equivalent to saying that $\sum_{i=1}^{\infty} i(i-1)\lambda_i = L''(1) > L'(1) = \sum_{i=1}^{\infty} i\lambda_i$, where $L(s)$ is the generating function of $\mathcal{D}$.

One of the main tools we use in the present work is the configuration model, which was introduced in different versions by Bender and Canfield in [6] and Bollobás in [9]. If $\mathbf{d}$ is a degree sequence on V_n , for some $n \in \mathbb{Z}^+$, we define the set of points $P = P(\mathbf{d})$ as $\{1 \times [d_1], \dots, n \times [d_n]\}$, where $[d_i] = \{1, \dots, d_i\}$ if $d_i > 0$ or the empty set otherwise. That is to every vertex in V_n correspond d_i points. Clearly, there are $2M$ points in P . Thus observe that there are $\frac{(2M)!}{M!2^M}$ perfect matchings on P . If $M(\mathbf{d})$ is such a perfect matching, then we can obtain a (multi)graph $GM(\mathbf{d})$ if we project P onto V_n preserving adjacencies, namely for any two vertices $i, j \in V_n$, if $M(\mathbf{d})$ contains an edge between a point in $i \times [d_i]$ and a point in $j \times [d_j]$, then $GM(\mathbf{d})$ contains a copy of the edge (i, j) . Of course in a perfect matching there might be edges that join two points corresponding to the same vertex, in which case $GM(\mathbf{d})$ obtains a loop on the vertex. Similarly, there might be two vertices which are joined to each other with more than one pairs of points, and in this case $GM(\mathbf{d})$ obtains multiple copies of the corresponding edge. If $M(\mathbf{d})$ is a uniformly random perfect matching on P , then observe that $GM(\mathbf{d})$ is not uniformly distributed over the set of multigraphs having $\mathbf{d}$ as their degree sequence. However, if we condition on the event that $GM(\mathbf{d})$ is a simple graph, then it is uniformly distributed over the set of simple graphs that have $\mathbf{d}$ as their degree sequence.

Consider now an asymptotic degree sequence $\mathcal{D} = (\mathbf{d}(n))_{n \in \mathbb{Z}^+}$. For each $n \in \mathbb{Z}^+$ we set $P(n)$ to be the set of points that corresponds to the degree sequence $\mathbf{d}(n)$. Let M_n be a uniformly random perfect matching on $P(n)$ and let $\tilde{G}(n)$ be the multigraph that is obtained from the projection of M_n onto V_n . The following theorem was proved by M. Molloy and B. Reed in [28] and has a key role in our proofs:

Theorem 2.1 *Let $\mathcal{D} = (\mathbf{d}(n))_{n \in \mathbb{Z}^+}$ be a sparse asymptotic degree sequence of maximum degree at most $n^{1/9}$.*

- *If $Q(\mathcal{D}) > 0$, then there exists an $\varepsilon > 0$ such that $\mathbb{P}[|L_1(\tilde{G}(n))| \geq \varepsilon n] \rightarrow 1$, as $n \rightarrow \infty$.*
- *If $Q(\mathcal{D}) < 0$, then for every $\varepsilon > 0$ we have $\mathbb{P}[|L_1(\tilde{G}(n))| \geq \varepsilon n] \rightarrow 0$, as $n \rightarrow \infty$.*

Of course the above theorem as stated in [28] was referring to simple graphs rather than multigraphs. However, it was actually proved for the random multigraph $\tilde{G}(n)$ and conditioning on being simple it can be stated for random simple graphs having this particular degree sequence. In fact in the first case Molloy and Reed proved the uniqueness of the component that has linear order; in particular the second largest component has logarithmic order. The restriction on the maximum degree can be slightly relaxed (see [28]), but for the simplicity of our proofs we assume it to be as above. The way we use this result will become apparent during the sketch of our proofs that is about to follow.

Note that assuming that $\mathcal{D}$ has $L''(1) > L'(1)$, the above theorem implies that $\tilde{G}(n)$ will have a giant component with probability $1 - o(1)$.

Here are the two deletion processes that we consider separately:

- *bond percolation*: For some $p \in (0, 1)$, we delete at random each edge of $\tilde{G}(n)$ with probability $1 - p$, independently of every other edge.
- *site percolation*: with probability $1 - p$ we make a vertex isolated by deleting the edges that are incident to it, independently for every vertex of $\tilde{G}(n)$.

In either case, the random multigraph that is the outcome of this experiment is denoted by $G'(n)$.

Eventually we want to know the structure of $G'(n)$, if $\tilde{G}(n)$ is a simple graph. Thus, we will show that if $\mathcal{A}(n)$ is set of multigraphs on V_n and $\mathbb{P}[G'(n) \in \mathcal{A}(n)] \rightarrow 0$ as $n \rightarrow \infty$, then $\mathbb{P}[G'(n) \in \mathcal{A}(n) \mid \tilde{G}(n) \text{ is simple}] \rightarrow 0$ as $n \rightarrow \infty$ as well. Hence, it will be sufficient for our purposes to perform the random deletion on the edges or the vertices of a random perfect matching on $P(n)$ without any conditioning and henceforth to consider the multigraph that is obtained from the remaining edges; this is going to make the calculations much simpler.

Thus we prove the following lemma:

Lemma 2.2 *Let $\mathcal{A}(n)$ be a set of multigraphs on V_n and suppose that $\mathbb{P}[G'(n) \in \mathcal{A}(n)] \rightarrow 0$ as $n \rightarrow \infty$. Then $\lim_{n \rightarrow \infty} \mathbb{P}[G'(n) \in \mathcal{A}(n) \mid \tilde{G}(n) \text{ is simple}] = 0$ as well.*

Proof. Note that

$$\mathbb{P}[G'(n) \in \mathcal{A}(n) \mid \tilde{G}(n) \text{ is simple}] \leq \frac{\mathbb{P}[G'(n) \in \mathcal{A}(n)]}{\mathbb{P}[\tilde{G}(n) \text{ is simple}]} \quad (5)$$

The asymptotic enumeration formula for graphs with a given degree sequence, such that $M = \Theta(n)$ and $\Delta = o(n^{1/3})$, obtained by McKay and Wormald in [26] yields

$$\mathbb{P}[\tilde{G}(n) \text{ is simple}] = (1 + o(1))e^{-\lambda/2 - \lambda^2/4},$$

where $\lambda = \frac{1}{M} \sum_{i=1}^n \binom{d_i}{2}$. But the latter sum is at most $\sum_{i=1}^n d_i^2 = O(n)$, and since $M = \Omega(n)$ it follows that $\lambda = O(1)$. Thus

$$\liminf_{n \rightarrow \infty} \mathbb{P}[\tilde{G}(n) \text{ is simple}] > 0,$$

and this concludes the proof of the lemma as the numerator in (5) converges to zero. $\blacksquare$

In both cases the random deletion induces a (random) degree sequence on V_n which we denote by $\mathbf{d}'(n)$ (the use the same symbol for the two kinds of percolation should cause no confusion). So, for each $n \in \mathbb{Z}^+$, let $\mathbf{D}_n$ be the set of degree sequences on V_n that are the result of the random deletion equipped with the probability distribution inherited by the random experiment we just described. That is, the probability of a certain degree sequence $\mathbf{d}'(n) \in \mathbf{D}_n$ is the probability

that the degree sequence which is induced by the random deletion (either of edges or of vertices) on $\tilde{G}(n)$ is $\mathbf{d}'(n)$. We set $\mathbf{D} = \prod_{n=1}^{\infty} \mathbf{D}_n$ to be the product space equipped with the product measure, which we denote by μ . Thus each element of $\mathbf{D}$ is an asymptotic degree sequence and Q is now a random variable on $\mathbf{D}$.

The strategy of our proof is quite different from that in [23], in that we make explicit use of Theorem 2.1. We first prove that the perfect matching between those points that are the endpoints of the edges that survive the deletion either in bond percolation or in site percolation is uniformly random among the perfect matchings on these points. Hence to study the asymptotic properties of $G'(n)$ we shall condition first on its degree sequence for every $n \in \mathbb{Z}^+$ and then we shall study the asymptotic behaviour of $G'(n)$ conditioned on this asymptotic degree sequence. Of course to show that $G'(n)$ has a certain property with probability tending to 1 as $n \rightarrow \infty$, we have to show that almost all the asymptotic degree sequences in $\mathbf{D}$ have similar behaviour. In particular, if $D'_i(n)$ is the number of vertices of degree i in $G'(n)$, we shall prove that the random variable $\frac{1}{n} \sum_{i \geq 1} i(i-2)D'_i(n)$ converges μ -almost surely (μ -a.s.) to a quantity Q' that depends only on the λ_i 's and on p , which we will calculate explicitly in both cases. From this we derive the critical p_c , which we denote by p_c^{bond} for the case of bond percolation and p_c^{site} for the case of site percolation. We show that if $p > p_c$ then Q' is positive, whereas if $p < p_c$ we have $Q' < 0$. Using Theorem 2.1, we deduce the sudden appearance of a giant component in $G'(n)$ when p crosses p_c , with probability that tends to 1 as $n \rightarrow \infty$.

We conclude this section stating a concentration inequality which we use in our proofs and it follows from Theorem 7.1 in [27]. Let S be a finite set and let f be a real-valued function on the set of those subsets of S that have size k . Assume that whenever c, c' are two such subsets whose symmetric difference is 2, then $|f(c) - f(c')| \leq 2$. If C is chosen uniformly at random among the k -subsets of S , then

$$\mathbb{P}[|f(C) - \mathbb{E}[f(C)]| > t] \leq 2 \exp\left(-\frac{t^2}{2k}\right). \quad (6)$$

3 Bond percolation

In this case, we start with the random graph $\tilde{G}(n)$ which is the multigraph that is the projection onto V_n of a uniformly random perfect matching on $P(n)$ and we create the multigraph $G'(n)$, deleting each edge of the matching with probability $1 - p$, independently of every other edge. Thus, the number of edges of $G'(n)$ is distributed as $\text{Bin}(M(n), p)$.

Firstly, we will prove that the perfect matching on the remaining points in $P(n)$ conditional on the degree sequence that is created after the deletion is uniformly distributed on the set of perfect matchings on the set of points in $P(n)$ that survive the deletion. In particular, if C is the set points in $P(n)$ that are the end-points of the surviving edges, then for every $i \in V_n$ the new degree of vertex i is $|C \cap (i \times [\mathbf{d}_i(n)])|$. Hence the (random) set C induces a degree sequence on V_n , which we denote $\mathbf{d}'(n)$. We set $P'(n) = P(\mathbf{d}'(n))$.

Let $d \in \{0, \dots, \Delta\}$ and assume that the vertices $i_1, \dots, i_{k_d}$ (and only these) have new degree equal to d after the edge deletion. Hence $\mathbf{d}'(n)$ contains exactly k_d vertices of degree d and assume that these are $i, \dots, i + k_d - 1$. We identify i_j with $i + j - 1$, for every $j = 1, \dots, k_d$. Moreover, provided that $d \geq 1$ we also identify the d points of $C \cap (i_j \times [d(i_j)])$ with the points $\{i + j - 1\} \times \{1, \dots, d\}$ in $P'(n)$. Hence any perfect matching between the points in C corresponds to a perfect matching on $P'(n)$ and vice-versa. In other words, we obtain a bijection between the perfect matchings on these two sets of points.

In the case of bond percolation, the set $\mathbf{D}_n$ consists precisely of those degree sequences that are induced by the deletion of the edges of a random perfect matching on $P(n)$. The probability of a certain degree sequence in $\mathbf{D}_n$ will therefore be the probability that this is the induced degree sequence after the deletion. Our aim is to show that conditional on $\mathbf{d}'(n) = \mathbf{d}'$, each perfect matching on $P'(n) = P(\mathbf{d}')$ is equilikely.

To do so, we first prove the following:

Lemma 3.1 *Conditional on having k edges that survive the random deletion of the edges of the perfect matching on $P(n)$, the set of their $2k$ end-points is uniformly distributed among the $2k$ -subsets of $P(n)$.*

Proof. The probability that a specific $2k$ -subset of $P(n)$ is the set of the end-points of the k edges that survive is the probability that the perfect matching on $P(n)$ consists of a perfect matching on these $2k$ points and a perfect matching between the $2M - 2k$ remaining points, and that it is the set of the k edges on this $2k$ -subset that survive the deletion. The probability of this event is exactly:

$$\frac{\frac{(2M-2k)!}{(M-k)!2^{M-k}} \frac{2k!}{k!2^k}}{\frac{2M!}{M!2^M}} \frac{1}{\binom{M}{k}} = \frac{1}{\binom{2M}{2k}}.$$

This concludes the proof of the lemma. ■

With a little more work we obtain what we were aiming for:

Lemma 3.2 *Let $\mathbf{d}'(n)$ be the degree sequence that is induced by the random deletion of the edges of a uniformly random perfect matching on $P(n)$. Conditional on $\mathbf{d}'(n) = \mathbf{d}'$, any perfect matching on $P(\mathbf{d}')$ is equilikely.*

Proof. Assume that the sum of the degrees in $\mathbf{d}'$ is $2k$ and let $S_{\mathbf{d}'}$ be the set of those $2k$ -subsets of $P(n)$ that induce the degree sequence $\mathbf{d}'$. Let m be a particular perfect matching on $P'(n)$, conditional on $\mathbf{d}'(n) = \mathbf{d}'$. In other words, m is a perfect matching on $P(\mathbf{d}')$. Now, let us condition on $|P'(n)| = 2k$. If $C' \in S_{\mathbf{d}'}$, the probability that $C = C'$ and the particular perfect matching that corresponds to m is realised on C is $\frac{1}{\binom{M}{k}} \frac{\frac{(2M-2k)!}{(M-k)!2^{M-k}}}{\frac{2M!}{M!2^M}}$. Thus,

$$\mathbb{P}[m, \mathbf{d}'(n) = \mathbf{d}' \mid |P'(n)| = 2k] = \sum_{C' \in S_{\mathbf{d}'}} \frac{1}{\binom{M}{k}} \frac{\frac{(2M-2k)!}{(M-k)!2^{M-k}}}{\frac{2M!}{M!2^M}} = \frac{|S_{\mathbf{d}'}|}{\binom{M}{k}} \frac{\frac{(2M-2k)!}{(M-k)!2^{M-k}}}{\frac{2M!}{M!2^M}}. \quad (7)$$

By the previous lemma, conditional on $|P'(n)| = 2k$ every set in $S_{\mathbf{d}'}$ has probability $1/\binom{2M}{2k}$. Therefore,

$$\mathbb{P}[\mathbf{d}'(n) = \mathbf{d}' \mid |P'(n)| = 2k] = \frac{|S_{\mathbf{d}'}|}{\binom{2M}{2k}}.$$

Now, Bayes' rule (i.e. dividing (7) by the above probability) yields:

$$\begin{aligned} \mathbb{P}[m \mid \mathbf{d}'(n) = \mathbf{d}', |P'(n)| = 2k] &= \frac{1}{\binom{M}{k}} \frac{\frac{(2M-2k)!}{(M-k)!2^{M-k}}}{\frac{2M!}{M!2^M}} \binom{2M}{2k} \\ &= \frac{k!(M-k)!}{M!} \frac{\frac{(2M-2k)!}{(M-k)!2^{M-k}}}{\frac{2M!}{M!2^M}} \frac{2M!}{(2M-2k)!2k!} = \frac{1}{\frac{2k!}{k!2^k}}. \end{aligned}$$

But

$$\begin{aligned}
\mathbb{P}[m \mid \mathbf{d}'(n) = \mathbf{d}'] &= \mathbb{P}[m \cap |P'(n)| = 2k \mid \mathbf{d}'(n) = \mathbf{d}'] \\
&= \mathbb{P}[m \mid |P'(n)| = 2k, \mathbf{d}'(n) = \mathbf{d}'] \mathbb{P}[|P'(n)| = 2k \mid \mathbf{d}'(n) = \mathbf{d}'] \\
&= \mathbb{P}[m \mid |P'(n)| = 2k, \mathbf{d}'(n) = \mathbf{d}'],
\end{aligned}$$

since $\mathbb{P}[|P'(n)| = 2k \mid \mathbf{d}'(n) = \mathbf{d}'] = 1$, and the lemma follows. $\blacksquare$

For $i \in \mathbb{N}$, let $D'_i(n)$ be the number of vertices of degree i in $\mathbf{d}'(n)$. The threshold probability will be determined by the quantity $\sum_{i \geq 1} i(i-2)\lambda_i^{\text{bond}}$, where

$$\lambda_i^{\text{bond}} = \lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{E}[D'_i(n)].$$

Hence we need to determine each λ_i^{bond} , proving the existence of this limit, and to do so we will first calculate the expected value of $D'_i(n)$. We begin with the conditional expectation given the size of C , that is conditional on the number of points in $P(n)$ that survive the deletion. For any $k = 0, \dots, M$ we have

$$\mathbb{E}[D'_i(n) \mid |C| = 2k] = \sum_{d=i}^{\Delta} D_d(n) \mathbb{P}[\text{a given vertex of degree } d \text{ has new degree } i \mid |C| = 2k].$$

Recall that $|C|/2$, which equals the number of edges that survive the random deletion, is distributed as $\text{Bin}(M, p)$. Therefore, a standard concentration argument yields

$$\mathbb{P}[||C|/2 - Mp| > \ln n \sqrt{n}] \leq \exp(-\Omega(\ln^2 n)). \quad (8)$$

This indicates that we may restrict ourselves to $k \in I = [Mp - \ln n \sqrt{n}, Mp + \ln n \sqrt{n}]$.

By Lemma 3.1 conditional on $|C| = 2k$, the set C is uniformly distributed among all $2k$ -subsets of $P(n)$. Hence, we obtain

$$\begin{aligned}
&\mathbb{P}[\text{a given vertex of degree } d \text{ has new degree } i \mid |C| = 2k] = \\
&\binom{d}{i} \frac{\binom{2M-d}{2k-i}}{\binom{2M}{2k}} = \binom{d}{i} \frac{(2M-d)!}{2M!} \frac{2k!}{(2k-i)!} \frac{(2M-2k)!}{(2M-d-2k+i)!} \\
&= \binom{d}{i} \frac{(2k)^i}{(2M)^d} (2M-2k)^{d-i} \left(1 + O\left(\frac{1}{n^{7/9}}\right)\right) = \binom{d}{i} p^i (1-p)^{d-i} \left(1 + O\left(\frac{\ln n}{n^{7/18}}\right)\right),
\end{aligned}$$

uniformly for any $d \leq \Delta$ and any $k \in I$.

Therefore, since $D'_d(n) \leq n$ by (8) we obtain:

$$\begin{aligned}
\mathbb{E}[D'_i(n)] &= \sum_{k=0}^M \mathbb{E}[D'_i(n) \mid |C| = 2k] \mathbb{P}[|C| = 2k] \\
&= \sum_{k \in I} \sum_{d=i}^{\Delta} D_d(n) \binom{d}{i} p^i (1-p)^{d-i} \left(1 + O\left(\frac{\ln n}{n^{7/18}}\right)\right) \mathbb{P}[|C| = 2k] \\
&\quad + o\left(\frac{1}{n^3}\right) \\
&= \left(1 + O\left(\frac{\ln n}{n^{7/18}}\right)\right) \sum_{d=i}^{\Delta} D_d(n) \binom{d}{i} p^i (1-p)^{d-i} + o\left(\frac{1}{n^3}\right).
\end{aligned}$$

For every $\varepsilon > 0$, if i' and n are large enough

$$\frac{1}{n} \sum_{d=i'+1}^{\Delta} D_d(n) \binom{d}{i} p^i (1-p)^{d-i} \leq \frac{1}{n} \sum_{d=i'+1}^{\Delta} D_d(n) < \varepsilon.$$

Therefore,

$$\frac{1}{n} \sum_{d=i}^{i'} D_d(n) \binom{d}{i} p^i (1-p)^{d-i} \leq \frac{1}{n} \sum_{d=i}^{\Delta} D_d(n) \binom{d}{i} p^i (1-p)^{d-i} \leq \frac{1}{n} \sum_{d=i}^{i'} D_d(n) \binom{d}{i} p^i (1-p)^{d-i} + \varepsilon.$$

Taking limits on both sides we obtain:

$$\sum_{d=i}^{i'} \lambda_d \binom{d}{i} p^i (1-p)^{d-i} \leq \liminf_{n \rightarrow \infty} \frac{1}{n} \sum_{d=i}^{\Delta} D_d(n) \binom{d}{i} p^i (1-p)^{d-i}$$

and

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{d=i}^{\Delta} D_d(n) \binom{d}{i} p^i (1-p)^{d-i} \leq \sum_{d=i}^{i'} \lambda_d \binom{d}{i} p^i (1-p)^{d-i} + \varepsilon.$$

Letting $i' \rightarrow \infty$ and then $\varepsilon \rightarrow 0$, we obtain the value of λ_i^{bond} :

$$\lim_{n \rightarrow \infty} \frac{\mathbb{E}[D'_i(n)]}{n} = \sum_{d=i}^{\infty} \lambda_d \binom{d}{i} p^i (1-p)^{d-i} \equiv \lambda_i^{bond}. \quad (9)$$

We will show that the critical probability p_c^{bond} is equal to the root of the equation $Q' := \sum_{i=1}^{\infty} i(i-2)\lambda_i^{bond} = 0$, which we denote by $\hat{p}_{bond}$. Firstly, let us calculate $\hat{p}_{bond}$. We have

$$\begin{aligned} \sum_{i=1}^{\infty} i(i-2)\lambda_i^{bond} &= \sum_{i=1}^{\infty} i(i-2) \sum_{d=i}^{\infty} \lambda_d \binom{d}{i} p^i (1-p)^{d-i} = \sum_{d=1}^{\infty} \lambda_d \sum_{i=1}^d i(i-2) \binom{d}{i} p^i (1-p)^{d-i} \\ &= \sum_{d=1}^{\infty} \lambda_d (dp(1-p) + (dp)^2 - 2dp) = \sum_{d=1}^{\infty} \lambda_d (1-p + dp - 2) dp \\ &= \sum_{d=1}^{\infty} \lambda_d ((d-1)p - 1) dp = \sum_{d=1}^{\infty} \lambda_d d(d-1)p^2 - p \sum_{d=1}^{\infty} d\lambda_d \\ &= p \left(p \sum_{d=1}^{\infty} \lambda_d d(d-1) - \sum_{d=1}^{\infty} d\lambda_d \right). \end{aligned}$$

Therefore,

$$\hat{p}_{bond} := \frac{L'(1)}{L''(1)}.$$

We now let $Q'_n = \frac{1}{n} \sum_{i \geq 2} i(i-2)D'_i(n)$ and will show that $\lim_{n \rightarrow \infty} Q'_n$ exists $\mu - a.s.$ and it is equal to Q' . Hence, the sign of Q' determines the sign of $\lim_{n \rightarrow \infty} Q'_n$ for almost every asymptotic degree sequence in $\mathbf{D}$.

For notational convenience, we set $X_{i',n} = \frac{1}{n} \sum_{i \leq i'} i(i-2)D'_i(n)$. Clearly, $Q'_n \geq X_{i',n}$. On the other hand, (2) implies that for every $\varepsilon > 0$ there exists $i_0 = i_0(\varepsilon)$ such that whenever $i' > i_0$ for n sufficiently large

$$\frac{1}{n} \sum_{i > i'} i(i-2)D_i(n) < \varepsilon. \quad (10)$$

Since $D'_i(n) \leq D_i(n)$, for any $i' > i_0$ we have

$$Q'_n \leq X_{i',n} + \varepsilon.$$

We shall prove that for every such i' , $\mu - a.s.$

$$\lim_{n \rightarrow \infty} X_{i',n} = \sum_{i \leq i'} i(i-2) \lambda_i^{bond} =: Q'_{i'}. \quad (11)$$

In turn, this will imply that for every $i' > i_0$

$$Q'_{i'} \leq \liminf_{n \rightarrow \infty} Q'_n \leq \limsup_{n \rightarrow \infty} Q'_n \leq Q'_{i'} + \varepsilon, \quad \mu - a.s.$$

Now, letting $i' \rightarrow \infty$ yields

$$Q' \leq \liminf_{n \rightarrow \infty} Q'_n \leq \limsup_{n \rightarrow \infty} Q'_n \leq Q' + \varepsilon, \quad \mu - a.s.$$

Since the choice of ε is arbitrary, we may eventually deduce that:

$$\lim_{n \rightarrow \infty} Q'_n = Q', \quad \mu - a.s. \quad (12)$$

So let us focus on proving (11). This will follow, if we show that for every $\varepsilon > 0$

$$\sum_n \mathbb{P} [|X_{i',n} - Q'_{i'}| > \varepsilon] < \infty. \quad (13)$$

(See for example Lemma 6.8 in [31].) We will deduce the above inequality, proving that the summands are $o(1/n^3)$.

Thus, we continue with estimating

$$\mathbb{P} [|X_{i',n} - Q'_{i'}| > \varepsilon],$$

for some fixed $\varepsilon > 0$. Note that for n sufficiently large

$$|\mathbb{E}[X_{i',n}] - Q'_{i'}| \leq \frac{\varepsilon}{2}.$$

Thus

$$\mathbb{P} [|X_{i',n} - Q'_{i'}| > \varepsilon] \leq \mathbb{P} [|X_{i',n} - \mathbb{E}[X_{i',n}]| > \frac{\varepsilon}{2}].$$

If the latter is realised, then there exists $i \leq i'$ for which

$$\frac{1}{n} |D'_i(n) - \mathbb{E}[D'_i(n)]| > \frac{\varepsilon}{2 \sum_{i \leq i'} i(i-2)}.$$

Therefore, setting $\varepsilon' = \frac{\varepsilon}{2 \sum_{i \leq i'} i(i-2)}$ we have

$$\mathbb{P} [|X_{i',n} - \mathbb{E}[X_{i',n}]| > \frac{\varepsilon}{2}] \leq \sum_{i \leq i'} \mathbb{P} \left[\frac{1}{n} |D'_i(n) - \mathbb{E}[D'_i(n)]| > \varepsilon' \right].$$

We now show that each summand is $o(1/n^3)$. To do so, we will condition on the size of C . Recall that by Lemma 3.1 conditional on $|C| = 2k$, the set C is uniformly distributed among

the $2k$ -subsets of $P(n)$. Moreover, if we replace one of the points in C with another one that does not belong to C , then $D'_i(n)$ can change by at most 2. Therefore, applying (6) we obtain uniformly for any $k \in I$:

$$\begin{aligned} \mathbb{P} [|D'_i(n) - \mathbb{E}[D'_i(n)]| > \varepsilon' n \mid |C| = 2k] &\leq 2 \exp \left(-\frac{\varepsilon'^2 n^2}{4k} \right) \\ &\leq 2 \exp \left(-\frac{\varepsilon'^2 n^2}{4(Mp + \ln n \sqrt{n})} \right) = o \left(\frac{1}{n^3} \right). \end{aligned}$$

Therefore, by (8)

$$\begin{aligned} \mathbb{P} [|D'_i(n) - \mathbb{E}[D'_i(n)]| > \varepsilon' n] &= \\ \sum_{k \in I} \mathbb{P} [|D'_i(n) - \mathbb{E}[D'_i(n)]| > \varepsilon' n \mid |C| = 2k] \mathbb{P}[|C| = 2k] &+ o \left(\frac{1}{n^3} \right) = o \left(\frac{1}{n^3} \right), \end{aligned}$$

for every i .

Now that we have proved (12), we are ready to conclude the proof that $p_c^{bond} = \hat{p}_{bond}$. Let $E \subseteq \mathbf{D}$ be the event over which $\lim_{n \rightarrow \infty} Q'_n = Q'$; recall that $\mu(E) = 1$. Let $(\mathbf{d}'(n))_{n \in \mathbb{Z}^+} \in E$. If we condition on $\mathbf{d}'(n)$ being the degree sequence on $G'(n)$, then Lemma 3.2 implies that $G'(n)$ is the multigraph that arises as the projection of a uniformly random perfect matching on $P'(n)$ onto V_n .

If $p < \hat{p}_{bond}$ then $Q' < 0$. For an arbitrary $\varepsilon > 0$, we define $A(n)$ to be the set of multigraphs on V_n whose largest component has no more than εn vertices, for an arbitrary $\varepsilon \in (0, 1)$. In this case, Theorem 2.1 implies that $\lim_{n \rightarrow \infty} \mathbb{P}[G'(n) \in A(n) \mid D(G'(n)) = \mathbf{d}'(n)] = 1$, for every $\varepsilon > 0$.

On the other hand, if $p > \hat{p}_{bond}$ then $Q' > 0$. Again by Theorem 2.1, we deduce that there exists $\varepsilon > 0$ such that for any $(\mathbf{d}'(n))_{n \in \mathbb{Z}^+} \in E$, if we define $A(n)$ to be the set of multigraphs on V_n whose largest component has at least εn vertices, then $\lim_{n \rightarrow \infty} \mathbb{P}[G'(n) \in A(n) \mid D(G'(n)) = \mathbf{d}'(n)] = 1$.

However, in either case we want to know the limit of $\mathbb{P}[G'(n) \in A(n)]$ as $n \rightarrow \infty$ without conditioning on the degree sequence. If $\omega \in \mathbf{D}$, then we let $\pi_n(\omega)$ denote the projection of ω onto its n -th factor - recall that this is a degree sequence on V_n . Thus this probability can be expressed as follows:

$$\begin{aligned} \mathbb{P}[G'(n) \in A(n)] &= \sum_{\mathbf{d}'(n) \in \mathbf{D}_n} \mathbb{P}[G'(n) \in A(n) \mid D(G'(n)) = \mathbf{d}'(n)] \mathbb{P}[D(G'(n)) = \mathbf{d}'(n)] \\ &= \sum_{\mathbf{d}'(n) \in \mathbf{D}_n} \int_{\{\omega \in \mathbf{D} : \pi_n(\omega) = \mathbf{d}'(n)\}} \mathbb{P}[G'(n) \in A(n) \mid D(G'(n)) = \mathbf{d}'(n)] \mu(d\omega) \\ &= \int \mathbb{P}[G'(n) \in A(n) \mid D(G'(n)) = \pi_n(\omega)] \mu(d\omega). \end{aligned}$$

Since the integrand is bounded below by 0, applying Fatou's Lemma, we obtain:

$$\begin{aligned} \liminf_{n \rightarrow \infty} \mathbb{P}[G'(n) \in A(n)] &= \liminf_{n \rightarrow \infty} \int \mathbb{P}[G'(n) \in A(n) \mid D(G'(n)) = \pi_n(\omega)] \mu(d\omega) \\ &\geq \int \liminf_{n \rightarrow \infty} \mathbb{P}[G'(n) \in A(n) \mid D(G'(n)) = \pi_n(\omega)] \mu(d\omega) \\ &= \int_E \liminf_{n \rightarrow \infty} \mathbb{P}[G'(n) \in A(n) \mid D(G'(n)) = \pi_n(\omega)] \mu(d\omega) = \int_E \mu(d\omega) = 1. \end{aligned}$$

Now, applying the Reverse Fatou's Lemma (since the integrand is bounded above by 1), we have

$$\begin{aligned}
\limsup_{n \rightarrow \infty} \mathbb{P}[G'(n) \in A(n)] &= \limsup_{n \rightarrow \infty} \int \mathbb{P}[G'(n) \in A(n) \mid D(G'(n)) = \pi_n(\omega)] \mu(d\omega) \\
&\leq \int \limsup_{n \rightarrow \infty} \mathbb{P}[G'(n) \in A(n) \mid D(G'(n)) = \pi_n(\omega)] \mu(d\omega) \\
&= \int_E \limsup_{n \rightarrow \infty} \mathbb{P}[G'(n) \in A(n) \mid D(G'(n)) = \pi_n(\omega)] \mu(d\omega) = \int_E \mu(d\omega) = 1.
\end{aligned}$$

The last two inequalities along with Lemma 2.2 complete the proof of Theorem 1.1 for the bond percolation process.

4 Site percolation

In this section, we are dealing with site percolation, where for $p \in (0, 1)$ fixed, we make each vertex of $\tilde{G}(n)$ isolated with probability $1 - p$, independently of every other vertex, deleting all of the edges that are attached to it. We will be referring to this process as the *deletion* of the vertices. This process applied to $\tilde{G}(n)$ induces a random degree sequence on V_n , which, as in the previous section, we denote by $\mathbf{d}'(n)$. Now consider the effect of the deletion on the uniformly random perfect matching on $P(n)$: if a vertex is deleted, then the points of $P(n)$ that are the end-points of the edges attached to that vertex are deleted (i.e. we remove them from $P(n)$). Eventually, we are left with a set points of $P(n)$ that are the end-points of the remaining edges, and we denote it by C . Also, we let $P'(n) = P(\mathbf{d}'(n))$. As in the case of bond percolation, we establish a bijection between the set of perfect matchings between the points in C and the set of perfect matchings on $P'(n)$. In turn, this gives rise to a bijection between the set of perfect matchings on C and the set of perfect matchings on $P'(n)$.

In the present setting, the set $\mathbf{D}_n$ consists of the degree sequences that are induced by the random deletion of the vertices. The probability of a certain degree sequence in $\mathbf{D}_n$ will therefore be the probability that this is the induced degree sequence after the deletion.

We now argue that conditional on the choice of the points of C , each perfect matching on C has the same probability. The perfect matching which is realised after the deletion of the vertices is obtained in two independent stages: firstly the uniform perfect matching on $P(n)$ is realised and afterwards the random deletion of the vertices takes place. It is the independence that allows us to consider these two random experiments in reverse order. Thus, we choose first those vertices that will be deleted and then we realise the perfect matching on $P(n)$. Let $P_1(n)$ and $P_2(n)$ denote the sets of points corresponding to the deleted vertices and the vertices that remain, respectively. Let B be the subset of points in $P_2(n)$ that are matched with points in $P_1(n)$. Observe now that conditioning on the choice of $P_1(n)$ and B is equivalent to conditioning on the choice of C , as the disjoint union of B and C is $P_2(n)$. Under this conditioning each perfect matching on C has the same probability, since the number of perfect matchings on the remaining points is the same for every perfect matching on C . Thus if $|C| = 2k$, then each perfect matching on C has probability $k!2^k/(2k)!$.

For a degree sequence $\mathbf{d}'$, we let $S_{\mathbf{d}'}$ be the set of subsets of $P(n)$ which realise $\mathbf{d}'$. Assume that the sum of the degrees in $\mathbf{d}'$ is $2k$. Note that if $\mathbf{d}'(n) = \mathbf{d}'$, then $P'(n) = P(\mathbf{d}')$. Hence if m is a perfect matching on $P(\mathbf{d}')$, then

$$\mathbb{P}[m \mid \mathbf{d}'(n) = \mathbf{d}'] = \sum_{C' \in S_{\mathbf{d}'}} \mathbb{P}[m \mid C = C', \mathbf{d}'(n) = \mathbf{d}'] \mathbb{P}[C = C' \mid \mathbf{d}'(n) = \mathbf{d}']$$

$$= \frac{k!2^k}{(2k)!} \sum_{C' \in S_{\mathbf{d}'}} \mathbb{P}[C = C' \mid \mathbf{d}'(n) = \mathbf{d}'] = \frac{k!2^k}{(2k)!}.$$

The parameter p_c^{site} will be determined by $\mathbf{d}'(n)$. If $D'_i(n)$ denotes the number of vertices that have degree i in $\mathbf{d}'(n)$, then letting

$$\lambda_d^{site} = \lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{E}[D'_i(n)] \quad (14)$$

we shall prove that this limit exists for every $i \in \mathbb{N}$. In fact we show that

$$\lambda_d^{site} = p \lambda_d^{bond}. \quad (15)$$

This implies that

$$\sum_i i(i-2) \lambda_i^{site} = p \sum_i i(i-2) \lambda_i^{bond}. \quad (16)$$

Let $\hat{p}_{site}$ be the root of $\sum_i i(i-2) \lambda_i^{site} = 0$. We will show that $p_c^{site} = \hat{p}_{site}$. Then, by (16), we will deduce that $p_c^{site} = p_c^{bond}$.

We now prove the existence of the limit in (14). First of all we estimate the number of points in $P_2(n)$. Then we shall condition on a certain realisation of $P_2(n)$ and afterwards we shall condition on the size of B (i.e. on the size of C). From this we will be able to estimate $\mathbb{E}[D'_i(n)]$. Let $D''_d = D''_d(n)$ be the number of vertices of degree d that survive the deletion - therefore $\mathbb{E}[D''_d] = D_d p$. The total degree in $P_2(n)$ is $M_2 = \sum_{d=1}^{\Delta} d D''_d$ and the linearity of expectation yields $\mathbb{E}[M_2] = \sum_{d=1}^{\Delta} d D_d p = 2Mp$. As for every n the maximum degree in $\mathbf{d}(n)$ is no more than $n^{1/9}$ a bounded differences inequality (see for example Theorem 5.7 in [27]) yields

$$\mathbb{P}[|M_2 - \mathbb{E}[M_2]| > n^{2/3} \ln n] \leq \exp(-\Omega(\ln^2 n)). \quad (17)$$

Now note that if we condition on $|B| = b$, then any b -subset of $P_2(n)$ is equilikely to occur as the set B and it is the points of B that are deleted along with the points of $P_1(n)$.

Thus if $P_{i-d}(i)$ denotes the probability that after the random allocation of B a certain vertex in $P_2(n)$ of degree i loses $i-d$ points, thus becoming a vertex of degree d , the expected value of D'_d is

$$\mathbb{E}[D'_d] = \sum_{i=d}^{\Delta} \mathbb{E}[D''_i] P_{i-d}(i).$$

But for every $\varepsilon > 0$, there exists i_0 such that $\sum_{i > i_0} D_i \leq \varepsilon n$, for n sufficiently large. Since $D''_i(n) \leq D_i(n)$ we obtain

$$\sum_{i=d}^{i_0} \mathbb{E}[D''_i] P_{i-d}(i) \leq \mathbb{E}[D'_d] \leq \sum_{i=d}^{i_0} \mathbb{E}[D''_i] P_{i-d}(i) + \varepsilon n. \quad (18)$$

We now calculate $P_{i-d}(i)$, for $d \leq i \leq i_0$. Suppose that $M_2 = m_2$ and $|B| = b$. Then if $P_{i-d}(i, b, m_2)$ is the conditional probability that after the random choice of the b points in $P_2(n)$, which has m_2 points, a certain vertex of degree i loses $i-d$ points we have

$$P_{i-d}(i, b, m_2) = \binom{i}{i-d} \frac{\binom{m_2-i}{b-i+d}}{\binom{m_2}{b}} = \binom{i}{i-d} \frac{(m_2-i)!}{m_2!} \frac{b!}{(b-i+d)!} \frac{(m_2-b)!}{(m_2-b-d)!}.$$

We shall assume that for any n sufficiently large $b \in [2Mp(1-p) - n^{2/3} \ln^2 n, 2Mp(1-p) + n^{2/3} \ln^2 n]$. Indeed the following holds:

Lemma 4.1 *Conditional on $M_2 \in I' := [2Mp - n^{2/3} \ln n, 2Mp + n^{2/3} \ln n]$ we have $b \in I := [2Mp(1-p) - n^{2/3} \ln^2 n, 2Mp(1-p) + n^{2/3} \ln^2 n]$ with probability $1 - \exp(-\Omega(\ln^2 n))$.*

Proof. Assume that $M_2 = m_2$ for some $m_2 \in I'$. Therefore, $M_1 = 2M - m_2 \in [2M(1-p) - n^{2/3} \ln n, 2M(1-p) + n^{2/3} \ln n]$. We shall also condition on a particular realisation of the sets $P_1(n)$ and $P_2(n)$.

The probability that a certain point in $P_2(n)$ is adjacent to a point in $P_1(n)$ is $\frac{M_1}{2M-1} = (1-p)(1 + O(n^{-1/3} \ln n))$. Then $\mathbb{E}[b] = 2Mp(1-p) \pm O(n^{2/3} \ln n)$.

We now show that b is concentrated around its expected value, using Theorem 7.1 from [27]. We first describe here the more general setting on which this theorem applies and afterwards we will consider b .

Let W be a finite probability space that is also a metric space with its metric denoted by d . Assume that $P_0, \dots, P_s$ is a sequence of partitions on W , such that P_{i+1} is a refinement of P_i , P_0 is the partition consisting of only one part, that is W , and P_s is the partition where each part is an element of W . Assume that whenever $A, B \in P_{i+1}$ and $C \in P_i$ are such that $A, B \subseteq C$, then there is a bijection $\phi : A \rightarrow B$ such that $d(x, \phi(x)) \leq c_i$.

Now, let V be a uniformly random element of W and let $f : W \rightarrow \mathbb{R}$ be a function on W satisfying $|f(x) - f(y)| \leq d(x, y)$. Then

$$\mathbb{P}[|f(V) - \mathbb{E}[f(V)]| > t] \leq 2 \exp \left(-2 \frac{t^2}{\sum_{i=1}^s c_i^2} \right). \quad (19)$$

In our context the uniform space of all perfect matchings on $P(n)$ will play the role of W . Let $\mathbf{M}$ denote it. Its metric will be the symmetric difference of any two perfect matchings, regarded as sets. It is easy to see that this satisfies the properties a metric has by its definition. We shall consider a series of partitions on $\mathbf{M}$ denoted by $P_0, \dots, P_{M-1}$, where P_0 is $\mathbf{M}$ itself and each part of P_{M-1} will be a perfect matching in $\mathbf{M}$. To define the i -th partition, we define an ordering on the edges of each perfect matching. Consider first a linear ordering of all the points in $P(n)$. This induces a linear ordering on the edges of a perfect matching: if e_1 and e_2 are two edges, then $e_1 < e_2$ if the smallest point in e_1 is smaller than the smallest point in e_2 . Now a part of P_i consists of those perfect matchings whose i smallest edges are a particular set of i edges, provided that such a set of perfect matchings is non-empty. We call such an i -set of edges a *prefix*. Moreover, given a perfect matching, we call its i smallest edges its *i -prefix*.

Given such an i -set of edges, let C be the set of perfect matchings that have these i edges as their i -prefix. Now consider two $i+1$ -subsets that contain this i -set and are both prefixes. Suppose that e_A and e_B are the last edges on which they differ. Let A and B respectively denote the sets of perfect matchings that have these two $i+1$ -sets as their $i+1$ -prefixes.

There is a natural bijection $\phi : A \rightarrow B$ between them. Observe first that the smallest vertex in e_A and e_B is the same. In particular, let us assume that $e_A = (x, y_A)$ and $e_B = (x, y_B)$. If m is a matching in A , then $\phi(m)$ is the matching in B , where y_A is adjacent to the vertex that y_B was adjacent to in m ; every other edge remains unchanged. Note that the symmetric difference of m and $\phi(m)$ is 4. In other words, $c_i = 4$.

Now, we are ready to apply the concentration bound (19) to b . For any $m \in \mathbf{M}$, we let $b(m)$ be the number of edges between $P_1(n)$ and $P_2(n)$. Observe that for any two perfect matchings $m, m' \in \mathbf{M}$, always $|b(m) - b(m')|$ is no more than the size of the symmetric difference of m and m' . Thus applying (19) with $t = n^{2/3} \ln^2 n / 2$, the lemma follows, for n large enough. ■

Thus, uniformly for any $b \in I$ and any $m_2 \in I'$ we have:

$$P_{i-d}(i, b, m_2) = \binom{i}{i-d} \frac{b^{i-d} (m_2 - b)^d}{m_2^i} \left(1 + O\left(\frac{1}{n}\right) \right)$$

$$= \binom{i}{d} (1-p)^{i-d} p^d \left(1 + O\left(\frac{\ln^2 n}{n^{1/3}}\right) \right).$$

A standard concentration argument shows that uniformly for any $i \leq i_0$ we have

$$\mathbb{P}[|D_i''(n) - \mathbb{E}[D_i''(n)]| \geq \sqrt{n} \ln n] \leq \exp(-\Omega(\ln^2 n)). \quad (20)$$

Thus if we also set $I''(i) = [\max\{pD_i - \ln n \sqrt{n}, 0\}, pD_i + \ln n \sqrt{n}]$, we have

$$\mathbb{P}[b \notin I \text{ or } M_2 \notin I' \text{ or } D_i'' \notin I''(i), \text{ for some } i \leq i_0] = o(n^{-3}). \quad (21)$$

Therefore, the right-hand side of (18) becomes

$$\begin{aligned} \mathbb{E}[D'_d] &\leq \sum_{k \in I} \sum_{k' \in I'} \sum_{i=d}^{i_0} \sum_{k'' \in I''(i)} k'' P_{i-d}(i, k, k') \mathbb{P}[b = k, M_2 = k', D_i'' = k''] + \varepsilon n + o(n^{-2}) \\ &= \sum_{i=d}^{i_0} \sum_{k'' \in I''(i)} k'' \binom{i}{d} (1-p)^{i-d} p^d \mathbb{P}[D_i'' = k''] \left(1 + O\left(\frac{\ln^2 n}{n^{1/3}}\right) \right) + \varepsilon n + o(n^{-2}). \end{aligned}$$

But by (20), we have $\sum_{k'' \in I''(i)} k'' \mathbb{P}[D_i'' = k''] = \mathbb{E}[D_i''(n)] - o(n^{-2}) = D_i(n)p - o(n^{-2})$. Substituting this into the above expression, (18) now yields:

$$\mathbb{E}[D'_d] \leq p \sum_{i=d}^{i_0} D_i(n) \binom{i}{d} (1-p)^{i-d} p^d \left(1 + O\left(\frac{\ln^2 n}{n^{1/3}}\right) \right) + \varepsilon n + o(n^{-2})$$

and also repeating the above estimations,

$$\mathbb{E}[D'_d] \geq p \sum_{i=d}^{i_0} D_i(n) \binom{i}{d} (1-p)^{i-d} p^d \left(1 + O\left(\frac{\ln^2 n}{n^{1/3}}\right) \right) + o(n^{-2}).$$

Therefore,

$$p \sum_{i=d}^{i_0} \lambda_i \binom{i}{d} (1-p)^{i-d} p^d \leq \liminf_{n \rightarrow \infty} \frac{1}{n} \mathbb{E}[D'_d]$$

and

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \mathbb{E}[D'_d] \leq p \sum_{i=d}^{i_0} \lambda_i \binom{i}{d} (1-p)^{i-d} p^d + \varepsilon.$$

Letting $i_0 \rightarrow \infty$ and $\varepsilon \rightarrow 0$ we obtain

$$\lambda_d^{site} \equiv \lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{E}[D'_d] = p \sum_{i=d}^{\infty} \lambda_i \binom{i}{d} (1-p)^{i-d} p^d, \quad (22)$$

which yields (15) through (9).

Now we let $Q'_n = \frac{1}{n} \sum_{i \geq 2} i(i-2) D'_i(n)$. We will show that $\mu - a.s.$

$$\lim_{n \rightarrow \infty} Q'_n = \sum_{i \geq 1} i(i-2) \lambda_i^{site} \equiv Q'. \quad (23)$$

To prove this we argue as in the case of bond percolation: setting $X_{i',n} = \frac{1}{n} \sum_{i \leq i'} i(i-2)D'_i(n)$, for every $\varepsilon > 0$ and any i' large enough we have

$$X_{i',n} \leq Q'_n \leq X_{i',n} + \varepsilon,$$

if n is also large enough. (Obviously the first inequality holds for every i' and n .) Thus the existence of the $\mu - a.s.$ limit of Q'_n will be established once we show that for any i' $\mu - a.s.$ $\lim_{n \rightarrow \infty} X_{i',n} = \sum_{i \leq i'} i(i-2)\lambda_i^{site} \equiv Q'_{i'}$. We then let $i' \rightarrow \infty$ and $\varepsilon \rightarrow 0$ to deduce that $\mu - a.s.$ $\lim_{n \rightarrow \infty} Q'_n = Q'$.

The almost sure convergence of $X_{i',n}$ to $Q'_{i'}$ can be shown as in the case of bond percolation. In other words, we need to prove that the condition in (13) is satisfied in the present context. As before, we will show that for every $i \leq i'$ the random variable $D'_i(n)$ is sharply concentrated around its expected value: that is its tails converge to 0 exponentially fast. Recall that the total degree in $P_2(n)$ is denoted by M_2 .

Conditional on a certain realisation of $P_2(n)$, with $M_2 = m_2$ for some $m_2 \in I'$, and $|B| = b$ for some $b \in I$, the value of $D'_i(n)$ is determined by the random choice of the set B in $P_2(n)$. Note that D'_i can change by at most 2, if we replace one element of B by another one. Therefore we may apply (6):

$$\begin{aligned} \mathbb{P} [|D'_i(n) - \mathbb{E}[D'_i(n)]| > \ln n \sqrt{n} \mid |B| = b, P_2(n), |P_2(n)| = m_2] \leq \\ 4 \exp \left(- \frac{n \ln^2 n}{2(2Mp(1-p) + n^{2/3} \ln^2 n)} \right) = \exp(-\Omega(\ln^2 n)), \end{aligned}$$

uniformly for any $b \in I$ and $m_2 \in I'$. Hence, the above inequality along with (17) and Lemma (4.1) imply that

$$\mathbb{P} [|D'_i(n) - \mathbb{E}[D'_i(n)]| > \ln n \sqrt{n}] = o(n^{-3}).$$

Since $i \leq i'$ and i' is bounded, condition (13) is satisfied, and therefore, $\mu - a.s.$ $\lim_{n \rightarrow \infty} X_{i',n} = \sum_{i \leq i'} i(i-2)\lambda_i^{site} \equiv Q'_{i'}$. Now this concludes the proof of (23).

The proof that $p_c^{site} = \hat{p}_{site}$ is identical to that for $\hat{p}_{bond}$, and it is omitted.

References

- [1] M. Ajtai, J. Komlós and E. Szemerédi, Largest random component of the k -cube, *Combinatorica* **2** (1982), 1–7.
- [2] R. Albert and A.L. Barabási, Statistical Mechanics of complex networks, *Reviews of Modern Physics* **74** (2002), 47–97.
- [3] R. Albert, H. Jeong and A.L. Barabási, Error and attack tolerance of complex networks, *Nature* **406** (2000), 378–382.
- [4] N. Alon, I. Benjamini and A. Stacey, Percolation on finite graphs and isoperimetric inequalities, *Annals of Probability* **32** (2004), 1727–1745.
- [5] K.B. Atreya and P.E. Ney, *Branching Processes*, Springer, 1972.
- [6] E.A. Bender and E.R. Canfield, The asymptotic number of labelled graphs with given degree sequences. *J. Combin. Theory Ser. A* **24** (1978), 296–307.

- [7] M. Boguñá, R. Pastor-Satorras and A. Vespignani, Absence of epidemic threshold in scale-free networks with degree correlations, *Phys. Rev. Lett.* **90** (2003), 028701-1–4.
- [8] B. Bollobás, *Random Graphs*, Cambridge University Press, 2001.
- [9] B. Bollobás, A probabilistic proof of an asymptotic formula for the number of labelled regular graphs, *Europ. J. Combinatorics* **1** (1980), 311–316.
- [10] B. Bollobás, C. Borgs, J. Chayes and O. Riordan, Percolation on dense graph sequences, preprint (available at [arXiv:math.PR/0701346](https://arxiv.org/abs/math.PR/0701346)).
- [11] B. Bollobás, Y. Kohayakawa and T. Łuczak, The evolution of random subgraphs of the cube. *Random Structures and Algorithms* **3** (1992), 55–90.
- [12] B. Bollobás and O. Riordan, Mathematical results on scale-free random graphs, In *Handbook on Graphs and Networks*, Wiley-VCH, Berlin, 2002.
- [13] C. Borgs, J. Chayes, R. van der Hofstad, G. Slade and J. Spencer, The scaling window under the triangle condition, *Random Structures and Algorithms* **27** (2005), 137–184.
- [14] C. Borgs, J. Chayes, R. van der Hofstad, G. Slade and J. Spencer, The lace expansion and the triangle condition, *Annals of Probability* **33**(5) (2005), 1886–1944.
- [15] C. Borgs, J. Chayes, R. van der Hofstad, G. Slade and J. Spencer, The phase transition for the n -cube, *Combinatorica* **26** (2006), 395–410.
- [16] D.S. Callaway, M.E.J. Newman, S.H. Strogatz and D.J. Watts, Network robustness and fragility: Percolation on random graphs, *Phys. Rev. Lett.* **85** (2000), 5468–5471.
- [17] F. Chung and L. Lu, *Complex Graphs and Networks*, AMS, 2006.
- [18] R. Cohen, K. Erez, D. ben-Avraham and S. Havlin, Resilience of the Internet to random breakdowns, *Phys. Rev. Lett.* **85** (2000), 4626–4628.
- [19] S.N. Dorogotsev and J.F.F. Mendes, Evolution of networks, *Advances in Physics* **51** (2002), 1079–1187.
- [20] P. Erdős and A. Rényi, On the evolution of random graphs, *Publication of the Mathematical Institute of the Hungarian Academy of Sciences* **5** (1960), 17–61.
- [21] M. Faloutsos, P. Faloutsos and C. Faloutsos, On power-law relationships of the Internet topology, *Comp. Comm. Rev.* **29** (1999), 251–262.
- [22] A. Frieze, M. Krivelevich and R. Martin, The emergence of a giant component in random subgraphs of pseudo-random graphs. *Random Structures and Algorithms* **24** (2004), 42–50.
- [23] A. Goerdt, The giant component threshold for random regular graphs with edge faults, *Theor. Comput. Sci.* **259** (2001), 307–321.
- [24] G.R. Grimmett, *Percolation*, 2nd edn., Grundlehren der mathematischen Wissenschaften, vol. **321**, Springer-Verlag, Berlin, 1999.
- [25] S. Janson, T. Łuczak and A. Ruciński, *Random Graphs*, Wiley Interscience, 2000.

- [26] B.D. McKay and N.C. Wormald, Asymptotic enumeration by degree sequence of graphs with degrees $o(n^{1/2})$, *Combinatorica* **11** (1991), 369–382.
- [27] C.J.H. McDiarmid, On the method of bounded differences, In *Surveys in Combinatorics* (J. Siemons Ed.) Lond. Math. Soc. Lecture Notes, 1989, 148–188.
- [28] M. Molloy and B. Reed, A critical point for random graphs with given degree sequence, *Random Structures and Algorithms* **6** (1995), 161–179.
- [29] S. Nikolettseas, K. Palem, P. Spirakis and M. Yung, Vertex disjoint paths and multiconnectivity in random graphs: Secure network computing. In *Proceedings ICALP 1994* LNCS 820, 508–519.
- [30] S. Nikolettseas and P. Spirakis, Expansion properties of random regular graphs with edge faults. In *Proceedings STACS 1995* LNCS 900, 421–432.
- [31] V.V. Petrov, *Limit theorems of probability theory: sequences of independent random variables*, Clarendon Press, 1995.